

ON THE LOWEST TWO-SIDED CELL IN AFFINE WEYL GROUPS

JÉRÉMIE GUILHOT

ABSTRACT. Bremke and Xi determined the lowest two-sided cell for affine Weyl groups with unequal parameters and showed that it consists of at most $|W_0|$ left cells where W_0 is the associated finite Weyl group. We prove that this bound is exact. Previously, this was known in the equal parameter case and when the parameters were coming from a graph automorphism. Our argument uniformly works for any choice of parameters.

1. INTRODUCTION

This paper is concerned with the theory of Kazhdan-Lusztig cells in a Coxeter group W , following the general setting of Lusztig [7]. This involves a weight function L which is an integer-valued function on W such that $L(ww') = L(w) + L(w')$ whenever $\ell(ww') = \ell(w) + \ell(w')$ (ℓ is the usual length function on W). We shall only consider weight function such that $L(w) > 0$ for all $w \neq 1$. The case where $L = \ell$ is known as the equal parameter case.

The partition of W into cells is known to play an important role in the study of the representations of the corresponding Hecke algebra.

Here, we are primarily concerned with affine Weyl groups. Let W be an irreducible affine Weyl group, together with a weight function L . Let W_0 be the finite Weyl group associated to W .

In the equal parameter case, Shi [8] described the lowest two-sided cell c_0 with respect to the preorder $\leq_{LR}$ using the Lusztig a -function (see [7] for further details on the a -function) as follows

$$c_0 = \{w \in W \mid a(w) = \ell(w_0)\}$$

where w_0 is the longest element of W_0 . Then, he gave an upper bound for the number of left cells contained in c_0 , namely $|W_0|$. In [9], he showed that this bound is exact and he described the left cells in c_0 . His proof involved some deep properties of the Kazhdan-Lusztig polynomials in the equal parameter case, such as the positivity of the coefficients.

Using the positivity property, Lusztig ([6]) proved, in the equal parameter case, a number of results concerning the a -function, such as

$$(P4) \quad \text{if } z \leq_{LR} z' \text{ then } a(z) \geq a(z')$$

and

$$(P11) \quad \text{if } z \leq_{LR} z' \text{ and } a(z) = a(z') \text{ then } z \sim_{LR} z'.$$

Since we know that $a(z) \leq \ell(w_0)$ for any $z \in W$, this shows that c_0 is the lowest two-sided cell with respect to $\leq_{LR}$.

In the unequal parameter case, the positivity of the coefficients of the Kazhdan-Lusztig polynomials does not hold anymore. However, Bremke [2] and Xi [10] proved that the lowest two-sided cell c_0 can be described in the same way as in [8], using the general a -function. Let J be a subset of S and consider the corresponding parabolic subgroup $W_J = \langle J \rangle$. We denote by w_J the longest element in W_J . Let

$$\tilde{\nu} = \max_{J \subset S, W_J \simeq W_0} L(w_J).$$

Then, we have

$$c_0 = \{w \in W \mid a(w) = \tilde{\nu}\}.$$

Let $\mathbf{S}$ be the set which consists of all the subset J of S such that $W_J \simeq W_0$ and $L(w_J) = \tilde{\nu}$. Then we have the following alternative description of c_0 , and it is this description with which we work throughout the paper

$$c_0 = \{w \in W \mid w = x.w_J.y, \ x, y \in W, \ J \in \mathbf{S}\}$$

where for any $x, y, w \in W$ the notation $w = x.y$ means that $w = xy$ and $\ell(w) = \ell(x) + \ell(y)$ (and similarly $w = x.y.z$ for $w, x, y, z \in W$).

In [2], Bremke showed that c_0 contains at most $|W_0|$ left cells. She proved that this bound is exact when the parameters are coming from a graph automorphism and, again, the proof involved some deep properties of the Kazhdan-Lusztig polynomials in the equal parameter case.

In this paper, we will prove that $|W_0|$ is the exact bound for the number of left cells in c_0 for any choice of parameters; see Theorem 4.6. The method is based on a variation of the induction of left cells as in [3]. The main new ingredient is a geometric argument to find a “local” bound on the degree of the structure constants of the Hecke algebra of W with respect to the standard basis; see Theorem 2.4. Our proof uniformly works for any choice of parameters.

2. MULTIPLICATION OF THE STANDARD BASIS AND GEOMETRIC REALIZATION

In this section, we introduce the Hecke algebra of a Coxeter group with respect to a weight function. Then, we present a geometric realization of an affine Weyl group. Finally, using this geometric realization, we give a bound on the degree of the structure constants with respect to the standard basis.

2.1. Weight functions, Hecke algebras. In this section, (W, S) denotes an arbitrary Coxeter system. The basic reference is [7]. Let L be a weight function. In this paper, we will only consider the case where $L(w) > 0$ for all $w \neq 1$. A weight function is completely determined by its value on S and must only satisfy $L(s) = L(t)$ if s and t are conjugate.

Let $\mathcal{A} = \mathbb{Z}[v, v^{-1}]$ and $\mathcal{H}$ be the generic Iwahori-Hecke algebra associated to (W, S) with parameters $\{L(s) \mid s \in S\}$. $\mathcal{H}$ has an $\mathcal{A}$ -basis $\{T_w \mid w \in W\}$, called the standard basis, with multiplication given by

$$T_s T_w = \begin{cases} T_{sw}, & \text{if } sw > w, \\ T_{sw} + (v^{L(s)} - v^{-L(s)})T_w, & \text{if } sw < w, \end{cases}$$

(here, “ $<$ ” denotes the Bruhat order) where $s \in S$ and $w \in W$.

Let $x, y \in W$. We write

$$T_x T_y = \sum_{z \in W} f_{x,y,z} T_z$$

where $f_{x,y,z} \in \mathcal{A}$ are the structure constants with respect to the standard basis.

In this paper, we will be mainly interested in the case where W is an irreducible affine Weyl group (with corresponding Weyl group W_0). In that case, it is known that there is a global bound for the degrees of the structure constants $f_{x,y,z}$. Namely, set $\nu = \ell(w_0)$, $\tilde{\nu} = L(w_0)$, where w_0 is the longest element of W_0 , and $\xi_s = v^{L(s)} - v^{-L(s)}$ for $s \in S$. In [2], Bremke proved that

- (1) As a polynomial in ξ_s , $s \in S$, the degree of $f_{x,y,z}$ is at most ν
- (2) The degree of $f_{x,y,z}$ in v is at most $\tilde{\nu}$.

Our aim will be to find a “local” bound for the degrees of these polynomials, which depends on $x, y \in W$. For this purpose, we will work with a geometric realization of W , as described in the next section.

2.2. Geometric realization. In this section, we present a geometric realization of an affine Weyl group. The basic references are [2, 5, 10].

Let V be an euclidean space of finite dimension $r \geq 1$. Let Φ be an irreducible root system of rank r and $\check{\Phi} \subset V^*$ the dual root system. We denote the coroot corresponding to $\alpha \in \Phi$ by $\check{\alpha}$ and we write $\langle x, y \rangle$ for the value of $y \in V^*$ at $x \in V$. Fix a set of positive roots $\Phi^+ \subset \Phi$. Let W_0 be the Weyl group of Φ . For $\alpha \in \Phi^+$ and $n \in \mathbb{Z}$, we define a hyperplane

$$H_{\alpha,n} = \{x \in V \mid \langle x, \check{\alpha} \rangle = n\}.$$

Let

$$\mathcal{F} = \{H_{\alpha,n} \mid \alpha \in \Phi^+, n \in \mathbb{Z}\}.$$

Any $H \in \mathcal{F}$ defines an orthogonal reflection σ_H with fixed point set H . We denote by Ω the group generated by all these reflections, and we regard Ω as acting on the right on V . An alcove is a connected component of the set

$$V - \bigcup_{H \in \mathcal{F}} H.$$

Ω acts simply transitively on the set of alcoves X .

Let S be the set of Ω -orbits in the set of faces (codimension 1 facets) of alcoves. Then S consists of $r+1$ elements which can be represented as the $r+1$ faces of an alcove. If a face f is contained in the orbit $t \in S$, we say that f is of type t .

Let $s \in S$. We define an involution $A \rightarrow sA$ of X as follows. Let $A \in X$; then sA is the unique alcove distinct from A which shares with A a face of type s . The set of such maps generates a group of permutations of X which is a Coxeter group (W, S) . In our case, it is the affine Weyl group usually denoted $\tilde{W}_0$. We regard W as acting on the left on X . It acts simply transitively and commutes with the action of Ω .

Let L be a weight function on W . In [2], Bremke showed that if a hyperplane H in $\mathcal{F}$ supports faces of type $s, t \in S$ then s and t are conjugate in W which implies that $L(s) = L(t)$. Thus we can associate a weight $c_H \in \mathbb{Z}$ to $H \in \mathcal{F}$ such that $c_H = L(s)$ if H supports a face of type s .

Assume that W is not of type $\tilde{A}_1$ or $\tilde{C}_r$ ($r \geq 2$); Let H, H' be two parallel hyperplanes such that H supports a face of type s and H' a face of type s' . Then, Bremke [2] proved that s and s' are conjugate. In other words, if W is not of type $\tilde{A}_1$ or $\tilde{C}_r$ then any two parallel hyperplanes have the same weight.

In this paper, we will often have to distinguish the case where W is of type $\tilde{A}_1$ or $\tilde{C}_r$, because of this property.

In the case where W is of type $\tilde{C}_r$ with generators $s_1, \dots, s_{r+1}$ and W_0 is generated by $s_1, \dots, s_r$, by symmetry of the Dynkin diagram, we can assume that $c_{s_1} \geq c_{s_{r+1}}$. Similarly, if W is of type $\tilde{A}_1$ with generators s_1, s_2 and $W_0 = \langle s_1 \rangle$, we can assume that $c_{s_1} \geq c_{s_2}$.

For a 0-dimensional facet λ of an alcove, define

$$m(\lambda) = \sum_{H \in \mathcal{F}, \lambda \in H} c_H.$$

We say that λ is a special point if $m(\lambda)$ is maximal.

Let T be the set of all special points. For $\lambda \in T$, denote by W_λ the stabilizer of the set of alcoves containing λ in their closure with respect to the action of W on X . It is a maximal parabolic subgroup of W . Let $S_\lambda = S \cap W_\lambda$ and write w_λ for the longest element of W_λ . Note that, following [2], and with our convention for $\tilde{C}_r$ and $\tilde{A}_1$, $0 \in V$ is a special point. Moreover, if $\lambda = 0 \in V$, the definition of W_λ is consistent with the definition of W_0 given before.

Let λ be a special point, a quarter with vertex λ is a connected component of

$$V - \bigcup_{H, \lambda \in H} H.$$

It is an open simplicial cone. It has r walls.

Let $H = H_{\alpha, n} \in \mathcal{F}$. Then H divides $V - H$ into two half-spaces

$$V_H^+ = \{x \in V \mid \langle x, \check{\alpha} \rangle > n\},$$

$$V_H^- = \{x \in V \mid \langle x, \check{\alpha} \rangle < n\}.$$

Finally, let A_0 be the fundamental alcove defined by

$$A_0 = \{x \in V \mid 0 < \langle x, \check{\alpha} \rangle < 1 \text{ for all } \alpha \in \Phi^+\}.$$

Let $A \in X$, $w \in W$. It is well known that the length of w is the number of hyperplanes which separate A and wA .

2.3. Multiplication of the standard basis. Let (W, S) be an irreducible affine Weyl group associated to the Weyl group (W_0, S_0) . Recall that, for $x, y \in W$, we have

$$T_x T_y = \sum_{z \in W} f_{x, y, z} T_z.$$

After the preparations in 2.2, we will now be able to find a “local” bound for the degree of the polynomials $f_{x, y, z}$ which depends on $x, y \in W$.

For two alcoves $A, B \in X$, let

$$H(A, B) = \{H \in \mathcal{F} \mid H \text{ separates } A \text{ and } B\}.$$

Let $\overline{\mathcal{F}}$ be the set of directions of hyperplanes in $\mathcal{F}$. For $i \in \overline{\mathcal{F}}$, we denote by $\mathcal{F}_i$ the set of all hyperplanes $H \in \mathcal{F}$ of direction i . The connected components of

$$V - \bigcup_{H \in \mathcal{F}_i} H$$

are called “strip of direction i ”. We denote by $U_i(A)$ the unique strip of direction i which contains A , for $A \in X$. There exists a unique $\alpha \in \Phi^+$ and a unique $n \in \mathbb{Z}$

such that

$$U_i(A) = \{\mu \in V \mid n < \langle \mu, \check{\alpha} \rangle < n+1\},$$

in other words

$$U_i(A) = V_{H_{\alpha,n}}^+ \cap V_{H_{\alpha,n+1}}^-.$$

We say that $U_i(A)$ is defined by $H_{\alpha,n}$ and $H_{\alpha,n+1}$.

Note that our definition of strips is slightly different from the one in [2], where the strips were the connected components of

$$V - \bigcup_{\substack{H \in \mathcal{F}_i \\ c_H = c_i}} H \quad \text{where} \quad c_i = \max_{H \in \mathcal{F}, \overline{H} = i} c_H.$$

In fact, as noticed in Section 2.2, if W is not of type $\tilde{A}_1$ or $\tilde{C}_r$, then the two definitions are the same.

Let $\sigma \in \Omega$. Let $i, j \in \overline{\mathcal{F}}$ such that $\sigma(i) = j$. We have

$$(U_i(A))\sigma = U_j(A\sigma)$$

and the strip $U_j(A\sigma)$ is defined by the two hyperplanes $(H_{\alpha,n_i})\sigma$ and $(H_{\alpha,n_i+1})\sigma$.

Let $x, y \in W$; then we define

$$H_{x,y} = \{H \in \mathcal{F} \mid H \in H(A_0, yA_0) \cap H(yA_0, xyA_0)\},$$

$$I_{x,y} = \{i \in \overline{\mathcal{F}} \mid \exists H, \overline{H} = i, H \in H_{x,y}\}.$$

For $i \in I_{x,y}$, let

$$c_{x,y}(i) = \max_{H \in \mathcal{F}, \overline{H} = i, H \in H_{x,y}} c_H$$

and

$$c_{x,y} = \sum_{i \in I_{x,y}} c_{x,y}(i).$$

We are now ready to state the main result of this section.

Theorem 2.4. *Let $x, y \in W$ and*

$$T_x T_y = \sum_{z \in W} f_{x,y,z} T_z \quad \text{where } f_{x,y,z} \in \mathcal{A}.$$

Then, the degree of $f_{x,y,z}$ in v is at most $c_{x,y}$.

3. PROOF OF THEOREM 2.4

In order to prove Theorem 2.4, we will need the following lemmas.

Lemma 3.1. *Let $x, y \in W$ and $s \in S$ be such that $x < xs$ and $y < sy$. We have*

$$c_{xs,y} = c_{x,sy}.$$

Proof. Let H_s be the unique hyperplane which separates yA_0 and syA_0 . Since $x < xs$ and $y < sy$, one can see that

$$H(A_0, yA_0) \cup \{H_s\} = H(A_0, syA_0),$$

$$H(A_0, yA_0) \cap \{H_s\} = \emptyset,$$

and

$$H(syA_0, xsyA_0) \cup \{H_s\} = H(yA_0, xsyA_0),$$

$$H(syA_0, xsyA_0) \cap \{H_s\} = \emptyset.$$

Therefore we have

$$\begin{aligned}
H_{x, sy} &= H(A_0, syA_0) \cap H(syA_0, xsyA_0) \\
&= (H(A_0, yA_0) \cup \{H_s\}) \cap H(syA_0, xsyA_0) \\
&= (H(A_0, yA_0) \cap H(syA_0, xsyA_0)) \cup (\{H_s\} \cap H(syA_0, xsyA_0)) \\
&= H(A_0, yA_0) \cap H(syA_0, xsyA_0)
\end{aligned}$$

and

$$\begin{aligned}
H_{xs, y} &= H(yA_0, xsyA_0) \cap H(A_0, yA_0) \\
&= (H(syA_0, xsyA_0) \cup \{H_s\}) \cap H(A_0, yA_0) \\
&= (H(syA_0, xsyA_0) \cap H(A_0, yA_0)) \cup (\{H_s\} \cap H(A_0, yA_0)) \\
&= H(syA_0, xsyA_0) \cap H(A_0, yA_0) \\
&= H_{x, sy}.
\end{aligned}$$

Thus $c_{x, sy} = c_{xs, y}$. $\square$

Lemma 3.2. *Let $x, y \in W$ and $s \in S$ be such that $xs < x$ and $sy < y$. We have*

$$c_{xs, sy} \leq c_{x, y}.$$

Proof. Let H_s be the unique hyperplane which separate yA_0 and syA_0 . One can see that

$$H_{xs, sy} = H_{x, y} - \{H_s\}.$$

The result follows. $\square$

Lemma 3.3. *Let $x, y \in W$ and $s \in S$ be such that $xs < x$ and $sy < y$. Let H_s be the unique hyperplane which separates yA_0 and syA_0 . Then we have*

$$\overline{H_s} \notin I_{xs, y} \quad \text{and} \quad \overline{H_s} \in I_{x, y}.$$

Proof. We have

$$\begin{aligned}
sy < y &\implies H_s \in H(A_0, yA_0), \\
xs < x &\implies H_s \in H(yA_0, xsyA_0).
\end{aligned}$$

Thus $H_s \in H_{x, y}$ and $\overline{H_s} \in I_{x, y}$.

Let $\alpha_s \in \Phi^+$ and $n_s \in \mathbb{Z}$ be such that $H_s = H_{\alpha_s, n_s}$. Assume that $n_s \geq 1$ (the case where $n_s \leq 0$ is similar). Since $H_s \in H(A_0, yA_0)$ and yA_0 has a facet contained in H_s , we have

$$n_s < \langle \mu, \check{\alpha}_s \rangle < n_s + 1 \text{ for all } \mu \in yA_0.$$

Therefore, for all $m > n_s$, we have $H_{\alpha_s, m} \notin H(A_0, yA_0)$.

Now, since $xs < x$, we have

$$xsyA_0 \subset \{\mu \in V \mid n_s < \langle \mu, \check{\alpha}_s \rangle\}.$$

Therefore, for all $m \leq n_s$, we have $H_{\alpha_s, m} \notin H(yA_0, xsyA_0)$. Thus, there is no hyperplane parallel to H_s in $H_{xs, y}$, as required. $\square$

We keep the setting of the previous lemma. We denote by σ_s the reflection with fixed point set H_s . Assume that $I_{xs, y} \neq \emptyset$ and let $i \in I_{xs, y}$. Recall that $U_i(yA_0)$ is the unique strip of direction i which contains yA_0 . Since $i \in I_{xs, y}$ we have

$$A_0 \not\subset U_i(yA_0) \text{ and } xsyA_0 \not\subset U_i(yA_0).$$

One can see that one and only one of the hyperplanes which defines $U_i(yA_0)$ lies in $H_{xs,y}$. We denote by $H^{(i)}$ this hyperplane.

Let $H \in H_{xs,y}$. By the previous lemma we know that H is not parallel to H_s . Consider the 4 connected components of $V - \{H, H_s\}$. We denote by E_{A_0} , E_{yA_0} , E_{syA_0} and E_{xsyA_0} the connected component which contains, respectively, A_0 , yA_0 , syA_0 and $xsyA_0$. Assume that $(H)\sigma_s \neq H$. Then, we have either

$$(H)\sigma_s \cap E_{yA_0} \neq \emptyset \text{ and } (H)\sigma_s \cap E_{A_0} \neq \emptyset$$

or

$$(H)\sigma_s \cap E_{xsyA_0} \neq \emptyset \text{ and } (H)\sigma_s \cap E_{syA_0} \neq \emptyset.$$

Furthermore, in the first case, $(H)\sigma_s$ separates E_{xsyA_0} and E_{syA_0} , and, in the second case, $(H)\sigma_s$ separates E_{yA_0} and E_{A_0} . In particular, we have

$$\begin{aligned} (H)\sigma_s \cap E_{yA_0} \neq \emptyset &\implies (H)\sigma_s \in H(syA_0, xsyA_0) \\ (H)\sigma_s \cap E_{xsyA_0} \neq \emptyset &\implies (H)\sigma_s \in H(A_0, yA_0). \end{aligned}$$

Moreover, we see that

$$\begin{aligned} (H)\sigma_s \cap E_{yA_0} \neq \emptyset &\implies (H)\sigma_s \in H(syA_0, xsyA_0) \\ &\implies (H)\sigma_s \in H(yA_0, xsyA_0), \end{aligned}$$

since H_s is the only hyperplane in $H(yA_0, syA_0)$ and $(H)\sigma_s \neq H_s$.

We will say that $H \in H_{xs,y}$ is of *s-type 1* if $(H)\sigma_s \cap E_{yA_0} \neq \emptyset$ and of *s-type 2* if $(H)\sigma_s \cap E_{xsyA_0} \neq \emptyset$. To sum up, we have

- if H is of *s-type 1* then $(H)\sigma_s \in H(yA_0, xsyA_0)$;
- if H is of *s-type 2* then $(H)\sigma_s \in H(A_0, yA_0)$.

We illustrate this result in Figure 1. Note that if $H, H' \in H_{xs,y}$ are parallel, then they have the same type.




FIGURE 1. *s*-type 1 and *s*-type 2 hyperplanes

Lemma 3.4. *Let $x, y \in W$ and $s \in S$ be such that $xs < x$ and $sy < y$. Let H_s be the unique hyperplane which separates yA_0 and syA_0 and let σ_s be the corresponding reflection. The following holds.*

a) *Let $H \in \mathcal{F}$. We have*

$$H \in H(yA_0, xsyA_0) \Rightarrow (H)\sigma_s \in H(yA_0, xyA_0).$$

- b) Let $H \in H_{xs,y}$ be of s -type 1; then $H \in H_{x,y}$.
- c) Let $H \in H_{xs,y}$ be of s -type 2; then $(H)\sigma_s \in H_{x,y}$.
- d) Let $H \in H_{xs,y}$ such that $(H)\sigma_s = H$; then $H \in H_{x,y}$.

Proof. We prove (a). Let $H \in H(yA_0, xsyA_0)$. Then $(H)\sigma_s$ separates $yA_0\sigma_s$ and $xsyA_0\sigma_s$. But we have

$$yA_0\sigma_s = syA_0 \quad \text{and} \quad xsyA_0\sigma_s = xssyA_0 = xyA_0.$$

Since $H \neq H_s$, we have $(H)\sigma_s \neq H_s$ and this implies that $(H)\sigma_s$ separates yA_0 and xyA_0 .

We prove (b). We have $H \in H_{xs,y} = H(A_0, yA_0) \cap H(yA_0, xsyA_0)$. The hyperplane H is of s -type 1 thus $(H)\sigma_s \in H(yA_0, xsyA_0)$. Using (a) we see that $H \in H(yA_0, xyA_0)$. Therefore, $H \in H_{x,y}$.

We prove (c). Since H is of s -type 2 we have $(H)\sigma_s \in H(A_0, yA_0)$. Moreover, $H \in H(yA_0, xsyA_0)$ thus, using (a), we see that $(H)\sigma_s \in H(yA_0, xyA_0)$. Therefore, $(H)\sigma_s \in H_{x,y}$.

We prove (d). Using (a), we see that $(H)\sigma_s = H \in H(yA_0, xyA_0)$ and since $H \in H_{xs,y} \subset H(A_0, yA_0)$, we get $H \in H_{x,y}$. □

Lemma 3.5. *Let $x, y \in W$ and $s \in S$ be such that $xs < x$ and $sy < y$. Let H_s be the unique hyperplane which separates yA_0 and syA_0 . There is an injective map φ from $I_{xs,y}$ to $I_{x,y} - \{\overline{H_s}\}$.*

Proof. Let σ_s be the reflection with fixed point set H_s . If $I_{xs,y} = \emptyset$ then the result is clear. We assume that $I_{xs,y} \neq \emptyset$. We define φ as follows.

- (1) If $(H^{(i)})\sigma_s \in H(A_0, yA_0)$ then set $\varphi(i) = \sigma_s(i)$;
- (2) set $\varphi(i) = i$ otherwise.

We need to show that $\varphi(i) \in I_{x,y} - \{\overline{H_s}\}$. The fact that $\varphi(i) \neq \overline{H_s}$ is a consequence of Lemma 3.3, where we have seen that $\overline{H_s} \notin I_{xs,y}$. Indeed, since $\varphi(i)$ is either i or $\sigma_s(i)$ and $i \neq \overline{H_s}$ we cannot have $\varphi(i) = \overline{H_s}$.

Let $i \in I_{xs,y}$ be such that $\sigma_s(H^{(i)}) \in H(A_0, yA_0)$. By Lemma 3.4 (a), we have $\sigma_s(H^{(i)}) \in H(yA_0, xsyA_0)$. It follows that $\sigma_s(H^{(i)}) \in H_{x,y}$ and $\sigma_s(i) \in I_{x,y}$ as required.

Let $i \in I_{xs,y}$ be such that $\sigma_s(H^{(i)}) \notin H(A_0, yA_0)$. Then $H^{(i)}$ is of s -type 1. By the previous lemma we have $H^{(i)} \in H_{x,y}$ and $i \in I_{x,y}$.

We show that φ is injective. Let $i \in I_{xs,y}$ be such that $\varphi(i) = \sigma_s(i)$ and assume that $\sigma_s(i) \in I_{xs,y}$. We have

$$(U_i(yA_0))\sigma_s = U_{\sigma_s(i)}(syA_0) = U_{\sigma_s(i)}(yA_0)$$

and $(H^{(i)})\sigma_s$ is one of the hyperplane which defines $U_{\sigma_s(i)}(yA_0)$. Furthermore since $(H^{(i)})\sigma_s \in H(A_0, yA_0)$ we must have $(H^{(i)})\sigma_s = H^{(\sigma_s(i))}$. It follows that $(H^{(\sigma_s(i))})\sigma_s \in H(A_0, yA_0)$ and $\varphi(\sigma_s(i)) = i$. The result follows. □

Lemma 3.6. *Let $x, y \in W$ and $s \in S$ such that $xs < x$ and $sy < y$. Let H_s be the unique hyperplane which separates yA_0 and syA_0 . We have*

$$c_{xs,y} \leq c_{x,y} - c_{x,y}(\overline{H_s}).$$

Proof. Let φ be as in the proof of the previous lemma. We keep the same notation. If $I_{xs,y} = \emptyset$ then the result is clear, thus we may assume that $I_{xs,y} \neq \emptyset$.

First assume that W is not of type $\tilde{C}_r$ ($r \geq 2$) or $\tilde{A}_1$. Then any two parallel hyperplanes have the same weight, therefore we obtain, for $i \in I_{x,y}$


$$c_{x,y}(i) = c_{H(i)}.$$

Moreover, since $c_H = c_{(H)\sigma}$ for all $H \in \mathcal{F}$ and $\sigma \in \Omega$, one can see that

$$c_{x,y}(i) = c_{x,y}(\varphi(i)),$$

and the result follows using Lemma 3.5.

Now, assume that W is of type $\tilde{C}_r$, with graph and weight function given by



In [2], Bremke proved that the only case where two parallel hyperplanes H, H' do not have the same weight is when one of them, say H , supports a face of type s_1 and H' supports a face of type s_{r+1} .

If $a = b$, then parallel hyperplanes have the same weight and we can conclude as before.

Now assume that $a > b$. Let $i \in \overline{\mathcal{F}}$ be such that not all the hyperplanes with direction i have the same weight. Let $H = H_{\alpha,n}$ be a hyperplane with direction i and weight a . Then $H_{\alpha,n-1}$ and $H_{\alpha,n+1}$ have weight b because otherwise all the hyperplanes with direction i would have weight a .

Let $i \in I_{x,y}$. If no hyperplane of direction i supports a face of type s_1 or s_{r+1} then, as before, we can conclude that

$$c_{x,y}(i) = c_{x,y}(\varphi(i)).$$

Now, in order to prove that $c_{x,y} \leq c_{x,y} - c_{x,y}(\overline{H_s})$, the only problem which may appear is when there exists $i \in I_{x,y}$ such that

$$c_{x,y}(i) = a \quad \text{and} \quad c_{x,y}(\varphi(i)) = b.$$

Fix such a $i \in I_{x,y}$. We claim that

- (1) $H^{(i)}$ is of s -type 1 and $(H^{(i)})\sigma_s \in H(A_0, yA_0)$;
- (2) $\sigma_s(i) \in I_{x,y}$, $\varphi(\sigma_s(i)) = i$ and

$$c_{x,y}(\sigma_s(i)) = b \quad \text{and} \quad c_{x,y}(i) = a.$$

We prove (1). Let $j \in I_{x,y}$ be such that $H^{(j)}$ is of s -type 2. Then $(H^{(j)})\sigma_s \in H(A_0, yA_0)$ and $\varphi(j) = \sigma_s(j)$. Let $H \in H_{x,y}$ be such that $\overline{H} = j$. Then H is also of s -type 2 and $(H)\sigma_s \in H_{x,y}$ (see Lemma 3.4(c)). It follows that $c_{x,y}(\varphi(j)) \geq c_{x,y}(j)$. Let $j \in I_{x,y}$ be such that $(H^{(j)})\sigma_s = H^{(j)}$. Then $(H^{(j)})\sigma_s \in H(A_0, yA_0)$ and $\varphi(j) = j$. Let $H \in H_{x,y}$ be such that $\overline{H} = j$. Then $(H)\sigma_s = H$ and $H \in H_{x,y}$ (see Lemma 3.4(d)). It follows that $c_{x,y}(\varphi(j)) \geq c_{x,y}(j)$.

Finally let $j \in I_{x,y}$ be such that $H^{(j)}$ is of s -type 1 and $H^{(j)} \notin H(A_0, yA_0)$. Then $\varphi(j) = j$. Let $H \in H_{x,y}$ be such that $\overline{H} = j$. Then H is also of s -type 1 and $H \in H_{x,y}$ (see Lemma 3.4(b)). It follows that $c_{x,y}(\varphi(j)) \geq c_{x,y}(j)$.

Thus since $c_{x,y}(\varphi(i)) < c_{x,y}(i)$, we get that $H^{(i)}$ is of s -type 1 and $(H^{(i)})\sigma_s \in H(A_0, yA_0)$.

We prove (2). We know that $H^{(i)}$ is of s -type 1 and $(H^{(i)})\sigma_s \in H(A_0, yA_0)$. Thus $(H^{(i)})\sigma_s \in H_{x,y}$ and $\varphi(i) = \sigma_s(i)$. In particular, since $c_{x,y}(\varphi(i)) = b$, we must

have $c_{(H^{(i)})\sigma_s} = b$, which implies that $c_{H^{(i)}} = b$.

Since $H^{(i)}$ is of s -type 1 we have $(H^{(i)})\sigma_s \in H(yA_0, xsyA_0)$ which implies that $(H^{(i)})\sigma_s \in H_{x,y}$. Thus $\sigma_s(i) \in I_{x,y}$. Arguing as in the proof of Lemma 3.5, we obtain $(H^{(i)})\sigma_s = H^{(\sigma_s(i))}$ and $\varphi(\sigma_s(i)) = i$.

Let $\alpha \in \Phi^+$ and $n \in \mathbb{Z}$ be such that $H^{(i)} = H_{\alpha,n}$. Since $c_{x,y}(i) = a$, one can see that one of the hyperplanes $H_{\alpha,n-1}$, $H_{\alpha,n+1}$ lies in $H_{x,y}$. We denote this hyperplane by H . Note that $c_H = a$ thus, since $c_{x,y}(\sigma_s(i)) = b$, we cannot have $(H)\sigma_s \in H_{x,y}$. Both hyperplanes $(H)\sigma_s$ and $(H^{(i)})\sigma_s$ separate yA_0 and xyA_0 but only $(H^{(i)})\sigma_s$ lies in $H_{x,y}$. This implies that A_0 lies in the strip defined by $(H)\sigma_s$ and $(H^{(i)})\sigma_s$. Since $(H^{(i)})\sigma_s = H^{(\sigma_s(i))}$ this shows that the only hyperplane of direction $\sigma_s(i)$ which lies in $H_{x,y}$ is $H^{(\sigma_s(i))}$. Thus we have $c_{x,y}(\sigma_s(i)) = b$. Moreover $\varphi(\sigma_s(i)) = i$ and H is of s -type 1, thus $H \in H_{x,y}$ (see Lemma 3.4 (b)) and $c_{x,y}(i) = a$, as required.

Let $I_>$ be the subset of $I_{x,y}$ which consists of the directions i such that $c_{x,y}(i) = a$ and $c_{x,y}(\varphi(i)) = b$. Using (1) and (2), we see that the set $\sigma_s(I_>)$ is a subset of $I_{x,y}$ such that for all $i \in \sigma_s(I_>)$ we have $c_{x,y}(i) = b$ and $c_{x,y}(\varphi(i)) = a$. Therefore we can conclude that $c_{x,y} \leq c_{x,y} - c_{x,y}(\overline{H_s})$ in the case where W is of type $\tilde{C}_r$ ($r \geq 2$).

In the case where W is of type $\tilde{A}_1$, the result is clear, since we always have $I_{x,y} = \emptyset$. The lemma is proved. $\square$

Proof of Theorem 2.4. Let $x, y \in W$ and

$$T_x T_y = \sum_{z \in W} f_{x,y,z} T_z \quad \text{where } f_{x,y,z} \in \mathcal{A}.$$

We want to prove that the degree of $f_{x,y,z}$ in v is less than or equal to $c_{x,y}$. We proceed by induction $\ell(x) + \ell(y)$.

If $\ell(x) + \ell(y) = 0$ the result is clear.

If $c_{x,y} = 0$ then $H_{x,y} = \emptyset$ and $xy = x.y$. Thus $T_x T_y = T_{xy}$ and the result follows.

We may assume that $H_{x,y} \neq \emptyset$, which implies that $\ell(x) > 0$ and $\ell(y) > 0$. Let $x = s_k \dots s_1$ be a reduced expression of x . There exists $1 \leq i \leq k$ such that

$$\ell(s_{i-1} \dots s_1 y) = \ell(y) + i - 1 \text{ and } s_i s_{i-1} \dots s_1 y < s_{i-1} \dots s_1 y.$$

Let $x_0 = s_k \dots s_i$ and $y_0 = s_{i-1} \dots s_1 y$. Let H_{s_i} be the unique hyperplane which separates $y_0 A_0$ and $s_i y_0 A_0$. Note that $c_{H_{s_i}} = L(s_i)$. We have

$$T_x T_y = T_{x_0} T_{y_0}$$

Using Lemma 3.1, we obtain $c_{x,y} = c_{x_0,y_0}$. We have

$$\begin{aligned} T_{x_0,y_0} &= T_{s_k \dots s_{i+1}} T_{s_i} T_{y_0} \\ &= T_{s_k \dots s_{i+1}} (T_{s_i y_0} + \xi_{s_i} T_{y_0}) \\ &= T_{s_k \dots s_{i+1}} T_{s_i y_0} + \xi_{s_i} T_{s_k \dots s_{i+1}} T_{y_0} \\ &= T_{x_0 s_i} T_{s_i y_0} + \xi_{s_i} T_{x_0 s_i} T_{y_0} \end{aligned}$$

By induction, $T_{x_0 s_i} T_{s_i y_0}$ is an $\mathcal{A}$ -linear combination of T_z with coefficients of degree less than or equal to $c_{x_0 s_i, s_i y_0}$. Using Lemma 3.2, we have $c_{x_0 s_i, s_i y_0} \leq c_{x_0, y_0} = c_{x,y}$. By induction, $T_{x_0 s_i} T_{y_0}$ is an $\mathcal{A}$ -linear combination of T_z with coefficients of degree less than or equal to $c_{x_0 s_i, y_0}$. Therefore the degree of the polynomials occurring in

$\xi_{s_i} T_{x_0 s_i} T_{y_0}$ is less than or equal to $L(s_i) + c_{x_0 s_i, y_0}$. Applying Lemma 3.6 to x_0 and y_0 we obtain

$$c_{x_0 s_i, y_0} \leq c_{x_0, y_0} - c_{x_0, y_0}(\overline{H_{s_i}})$$

Since $c_{x_0, y_0}(\overline{H_{s_i}}) \geq c_{H_{s_i}} = L(s_i)$ we obtain

$$L(s_i) + c_{x_0 s_i, y_0} \leq c_{x_0, y_0} = c_{x, y}.$$

The theorem is proved. $\square$

4. THE LOWEST TWO-SIDED CELL

4.1. Kazhdan-Lusztig cells. Let (W, S) be a Coxeter group and L a weight function on W . Let $\mathcal{A} = \mathbb{Z}[v, v^{-1}]$ and $\mathcal{H}$ be the generic Iwahori-Hecke algebra corresponding to (W, S) with parameters $\{L(s) | s \in S\}$.

Let $a \mapsto \bar{a}$ be the involution of $\mathcal{A}$ which takes v^n to v^{-n} for all $n \in \mathbb{Z}$. We can extend it to a ring involution from $\mathcal{H}$ to itself by the formula

$$\overline{\sum_{w \in W} a_w T_w} = \sum_{w \in W} \bar{a}_w T_w^{-1}, \text{ where } a_w \in \mathcal{A}.$$

Let $\mathcal{A}_{\leq 0} = \mathbb{Z}[v^{-1}]$ and $\mathcal{A}_{< 0} = v^{-1}\mathbb{Z}[v^{-1}]$. For $w \in W$ there exists a unique element $C_w \in \mathcal{H}$ such that

$$\bar{C}_w = C_w \text{ and } C_w = T_w + \sum_{\substack{y \in W \\ y < w}} P_{y, w} T_y,$$

where $P_{y, w} \in \mathcal{A}_{< 0}$ for $y < w$. In fact, the set $\{C_w, w \in W\}$ forms a basis of $\mathcal{H}$, known as the Kazhdan-Lusztig basis. The elements $P_{y, w}$ are called the Kazhdan-Lusztig polynomials. We set $P_{w, w} = 1$ for any $w \in W$.

The Kazhdan-Lusztig left preorder $\leq_L$ on W is the relation generated by

$$\left\{ \begin{array}{l} y \leq_L w \text{ if there exists some } s \in S \text{ such that } C_y \text{ appears with} \\ \text{a non-zero coefficient in } T_s C_w, \text{ expressed in the } C_w\text{-basis} \end{array} \right.$$

One can see that

$$\mathcal{H} C_w \subseteq \sum_{y \leq_L w} \mathcal{A} C_y \text{ for any } w \in W.$$

The equivalence relation associated to $\leq_L$ will be denoted by $\sim_L$ and the corresponding equivalence classes are called the left cells of W . Similarly, we define $\leq_R$, $\sim_R$ and right cells. We say that $x \leq_{LR} y$ if there exists a sequence

$$x = x_0, x_1, \dots, x_n = y$$

such that for all $1 \leq i \leq n$ we have $x_{i-1} \leq_L x_i$ or $x_{i-1} \leq_R x_i$. We write $\sim_{LR}$ for the associated equivalence relation and the equivalence classes are called two-sided cells. The preorder $\leq_{LR}$ induces a partial order on the two-sided cells of W .

4.2. The lowest two-sided cell. Let (W, S) be an irreducible affine Weyl group. In this section, we look at the set

$$c_0 = \{w \in W \mid w = z'.w_\lambda.z, z, z' \in W, \lambda \in T\}$$

where T is the set of special points (see Section 2). We show that c_0 is the lowest two-sided cell and we determine the decomposition of c_0 into left cells.

Recall that, for λ a special point, W_λ is the stabilizer in W of the set of alcoves containing λ in their closure, w_λ is the longest element of W_λ and $S_\lambda = S \cap W_\lambda$. In particular, we have

$$sw_\lambda < w_\lambda \text{ for any } s \in S_\lambda.$$

For $\lambda \in T$ and $z \in W$ such that $w_\lambda z = w_\lambda \cdot z$, we set

$$N_{\lambda,z} = \{w \in W \mid w = z' \cdot w_\lambda \cdot z, z' \in W\}.$$

In [2, Proposition 5.1], it is shown that $N_{\lambda,z}$ is included in a left cell.

For $\lambda \in T$, we set

$$M_\lambda = \{z \in W \mid w_\lambda z = w_\lambda \cdot z, sw_\lambda z \notin c_0 \text{ for all } s \in S_\lambda\}.$$

Following [9], we choose a set of representatives for the Ω -orbits on T and denote it by R . Then

$$c_0 = \bigcup_{\lambda \in R, z \in M_\lambda} N_{\lambda,z} \quad (\text{disjoint union})$$

It is known ([9]) that this is a union over $|W_0|$ terms.

We are now ready to state the main result of this paper.

Theorem 4.3. *Let $\lambda \in T$ and $z \in M_\lambda$. The set $N_{\lambda,z}$ is a union of left cells. Furthermore, for $y \in W$ and $w \in N_{\lambda,z}$, we have*

$$y \leq_L w \implies y \in N_{\lambda,z}.$$

The proof of this theorem will be given in the next section. We now discuss a number of consequences of Theorem 4.3.

Corollary 4.4. *Let $\lambda \in T$ and $z \in M_\lambda$. Then, the set $N_{\lambda,z}$ is a left cell.*

Proof. The set $N_{\lambda,z}$ is a union of left cells which is included in a left cell (see [2, Proposition 5.1]). Hence it is a left cell. $\square$

The next step is to prove the following.

Proposition 4.5. *The set c_0 is included in a two-sided cell.*

Proof. Let $R = \{\lambda_1, \dots, \lambda_n\}$ be a set of representatives for the Ω -orbits on T . For example, if W is of type $\tilde{G}_2$ we have $n = 1$ and if W is of type $\tilde{B}_n$ ($n \geq 3$) we have $n = 2$. Set

$$c_{\lambda_i} = \{w \in W \mid w = z' \cdot w_{\lambda_i} \cdot z, z, z' \in W\}$$

One can see that

$$c_0 = \bigcup_{i=1}^{i=n} c_{\lambda_i}$$

and for $1 \leq i \leq j \leq n$ we have $c_{\lambda_i} \cap c_{\lambda_j} \neq \emptyset$. Therefore to prove the proposition, it is enough to show that each of the set c_{λ_i} is included in a two-sided cell.

Fix $1 \leq i \leq n$. Let $w, w' \in c_{\lambda_i}$ and $z, z', y, y' \in W$ be such that $w = z' \cdot w_{\lambda_i} \cdot z$ and $w' = y' \cdot w_{\lambda_i} \cdot y$. Using [2, Proposition 5.1], together with its version for right cells, we obtain

$$z' w_{\lambda_i} z \sim_L w_{\lambda_i} z \sim_R w_{\lambda_i} y \sim_R y' w_{\lambda_i} y.$$

The result follows. $\square$

Finally, combining the previous results of Shi, Xi and Bremke with Theorem 4.3, we now obtain the following description of the lowest two-sided cell in complete generality.

Theorem 4.6. *Let W be an irreducible affine Weyl group with associated Weyl group W_0 . Let*

$$c_0 = \{w \in W \mid w = z'.w_\lambda.z, \ z, z' \in W, \lambda \in T\}$$

where T is the set of special points. We have

- (1) c_0 is a two-sided cell.
- (2) c_0 is the lowest two-sided cell for the partial order on the two-sided cell induced by the preorder $\leq_{LR}$.
- (3) c_0 contains exactly $|W_0|$ left cells.
- (4) The decomposition of c_0 into left cells is as follows

$$c_0 = \bigcup_{\lambda \in R, \ z \in M_\lambda} N_{\lambda, z}.$$

Proof. We have seen that c_0 is included in a two-sided cell. Let $w \in c_0$ and $y \in W$ such that $y \sim_{LR} w$. In particular we have $y \leq_{LR} w$. We may assume that $y \leq_L w$ or $y \leq_R w$. We know that

$$c_0 = \bigcup_{\lambda \in R, \ z \in M_\lambda} N_{\lambda, z}.$$

Thus $w \in N_{\lambda, z}$ for some $\lambda \in R$ and $z \in M_\lambda$. If $y \leq_L w$ then, using Theorem 4.3, we see that $y \in N_{\lambda, z}$ and thus $y \in c_0$. If $y \leq_R w$ then using [7, §8.1], we have $y^{-1} \leq_L w^{-1}$. But c_0 is stable by taking the inverse, thus, as before, we see that $y^{-1} \in c_0$ and $y \in c_0$. This implies that c_0 is a two-sided cell and that it is the lowest one with respect to $\leq_{LR}$.

By [9], we know that

$$c_0 = \bigcup_{\lambda \in R, \ z \in M_\lambda} N_{\lambda, z}$$

is a disjoint union over $|W_0|$ terms. By Corollary 4.4, the result follows. $\square$

5. PROOF OF THEOREM 4.3

We keep the setting of the previous section. For $\lambda \in T$ we denote by X_λ the set of minimal left coset representatives of W_λ in W , that is

$$X_\lambda := \{w \in W \mid \ell(ws) > \ell(w) \text{ for all } s \in S_\lambda\}.$$

One can easily check that

$$X_\lambda = \{z \in W \mid zw_\lambda = z.w_\lambda\} \quad \text{and} \quad X_\lambda^{-1} = \{z \in W \mid w_\lambda z = w_\lambda.z\}.$$

Let $\lambda \in T$ and $z \in X_\lambda^{-1}$. For $z' \in W$, we have the following equivalence

$$z'w_\lambda z = z'.w_\lambda.z \iff z' \in X_\lambda.$$

Indeed, if $z'w_\lambda z = z'.w_\lambda.z$ then we must have $z'w_\lambda = z'.w_\lambda$ and $z' \in X_\lambda$. Conversely, if $z' \in X_\lambda$ then since $z \in X_\lambda^{-1}$ we have $z'w_\lambda z = z'.w_\lambda.z$ (see [9, Lemma 3.2]). Therefore we see that

$$N_{\lambda, z} = \{w \in W \mid w = xw_\lambda z, x \in X_\lambda\}.$$

5.1. Preliminaries.

Lemma 5.2. *Let $\lambda \in T$, $z \in M_\lambda$, $y \in X_\lambda$ and $v_1 < w_\lambda z$. Then, $P_{v_1, w_\lambda z} T_y T_{v_1}$ is an $\mathcal{A}$ -linear combination of $T_{z'}$ ($z' \in W$) with coefficient in $\mathcal{A}_{<0}$.*

Proof. We can write uniquely $v_1 = w.v'$, where $w \in W_\lambda$ and $v' \in X_\lambda^{-1}$ (see [4, Proposition 2.1.1]). First, assume that $w = w_\lambda$. In that case, we have $yv_1 = y.v_1$ and $T_y T_{v_1} = T_{yv_1}$. Since $P_{v_1, w_\lambda z} \in \mathcal{A}_{<0}$ the result follows.

Next, assume that $w < w_\lambda$. Let $w_{v_1} \in W$ such that $w_{v_1} w = w_\lambda$. The Kazhdan-Lusztig polynomials satisfy the following relation (see [7, Theorem 6.6.c])

$$P_{x,w} = v^{-L(s)} P_{sx,w}, \text{ where } x < sx \text{ and } sw < w.$$

Therefore, one can see that

- (1) $P_{v_1, w_\lambda z} \in v^{-L(w_{v_1})} \mathcal{A}_{<0}$ if $w_\lambda.v' < w_\lambda z$,
- (2) $P_{v_1, w_\lambda z} = v^{-L(w_{v_1})}$ if $w_\lambda.v' = w_\lambda z$.

Thus, to prove the lemma, it is sufficient to show that the polynomials occurring in the expression of $T_y T_{v_1}$ in the standard basis are of degree less than or equal to $L(w_{v_1})$ in the first case and $L(w_{v_1}) - 1$ in the second case.

Using Theorem 2.4, we know that the degree of these polynomials is less than or equal to c_{y,v_1} (for the definition of c_{y,v_1} , see Section 2.3).

Let $w_{v_1} = s_n \dots s_{m+1}$ and $w = s_m \dots s_1$ be reduced expressions, and let H_i be the unique hyperplane which separates $s_{i-1} \dots s_1 v' A_0$ and $s_i \dots s_1 v' A_0$. Note that $c_{H_i} = L(s_i)$. Let λ' be the unique special point contained in the closure of $v' A_0$ and $w_\lambda v' A_0$ (note that $W_{\lambda'} = W_\lambda$). By definition of X_λ , $yv_1 A_0$ lies in the quarter $\mathcal{C}$ with vertex λ' which contains $v_1 A_0$.

Let $1 \leq i \leq m$. Let α_i and $k \in \mathbb{Z}$ such that $H_i = H_{\alpha_i, k}$. Assume that $k > 0$ (the case $k \leq 0$ is similar). We have $v_1 A_0 \in V_{H_i}^+$. Now, since λ' lies in the closure of $v_1 A_0$ and $\lambda' \in H_i$, one can see that

$$k < \langle \mu, \tilde{\alpha}_i \rangle < k + 1 \text{ for all } \mu \in v_1 A_0.$$

Moreover, $yv_1 A_0 \in \mathcal{C}$ implies that

$$k < \langle \mu, \tilde{\alpha}_i \rangle \text{ for all } \mu \in yv_1 A_0.$$

From there, we conclude that all the hyperplanes $H_{\alpha_i, l}$ with $l \leq k$ do not lie in $H(v_1 A_0, yv_1 A_0)$ and that all the hyperplanes $H_{\alpha_i, l}$ with $l > k$ do not lie in $H(A_0, v_1 A_0)$. Thus $\overline{H_i} \notin I_{y,v_1}$ and we have

$$I_{y,v_1} \subset \{\overline{H_{m+1}}, \dots, \overline{H_n}\},$$

which implies

$$c_{y,v_1} \leq \sum_{i=m+1}^{i=n} c_{y,v_1}(\overline{H_i}).$$

Now, if W is not of type $\tilde{C}_r$ or $\tilde{A}_1$ then any two parallel hyperplanes have same weight and we have

$$c_{y,v_1}(\overline{H_i}) = \begin{cases} 0 & \text{if } i \notin I_{y,v_1}, \\ L(s_i) & \text{otherwise.} \end{cases}$$

Thus

$$c_{y,v_1} \leq \sum_{i=m+1}^{i=n} L(s_i) = L(w_{v_1}),$$

as required in the first case.

Assume that W is of type $\tilde{C}_r$ or $\tilde{A}_1$. Then, one can see that, since λ' is a special point, we have for all $1 \leq i \leq n$, $c_{H_i} = c_{\overline{H_i}} = L(s_i)$ and we can conclude as before.

Assume that we are in case 2. Let $j \in \overline{\mathcal{F}}$. Recall that in [2], the author defined the strip of direction j as the connected component of

$$V - \bigcup_{\substack{H \in \mathcal{F}, \overline{H}=j \\ c_H=c_j}} H, \quad \text{where} \quad c_j = \max_{H, \overline{H}=j} c_H.$$

To avoid confusion, we will call them maximal strips of direction j . Let

$$\mathcal{U}(A) = \bigcup_{U \text{ maximal strip, } A \subset U} U.$$

Now, $v_1 = w.v' < w_\lambda v' = w_\lambda z$ with $z \in M_\lambda$. Recall that

$$M_\lambda = \{z \in W \mid w_\lambda z = w_\lambda \cdot z, \text{ } sw_\lambda z \notin c_0 \text{ for all } s \in S_\lambda\}$$

thus $v_1 = w.v' \notin c_0$. In [1], Bédard showed (in the equal parameter case) that the lowest two-sided cell c_0 can be described as follows

$$c_0 = \{w \in W \mid wA_0 \not\subset \mathcal{U}(A_0)\}.$$

In [2], Bremke proved that this presentation remains valid in the unequal parameter case. Therefore since $v_1 A_0 \notin c_0$, we have $v_1 A_0 \in \mathcal{U}(A_0)$ and there exists a maximal strip U which contains A_0 and $v_1 A_0$. Let $1 \leq k \leq n$ be such that U is of direction $\overline{H}_k$. For $1 \leq i \leq m$, the hyperplane H_i separates A_0 and $v_1 A_0$ and $c_{H_i} = c_{\overline{H}_i}$, thus we must have $k > m$.

If W is not of type $\tilde{C}_r$ or $\tilde{A}_1$, then our strips and the strips as defined in [2] are the same. Therefore, since A_0 and $v_1 A_0$ lie in U , we have $\overline{H}_k \notin I_{y,v_1}$ and

$$c_{y,v_1} \leq \sum_{\substack{i=m+1 \\ i \neq k}}^{i=n} c_{y,v_1}(\overline{H}_i) \leq \sum_{\substack{i=m+1 \\ i \neq k}}^{i=n} L(s_i) < L(w_{v_1}),$$

as required.

Assume that W is of type $\tilde{C}_r$ or $\tilde{A}_1$. First, if all the hyperplanes with direction $\overline{H}_k$ have same weight, then we have $\overline{H}_k \notin I_{y,v_1}$ and we can conclude as before. Assume not, then we must have $c_{H_k} = c_{\overline{H}_k}$ (since $\lambda' \in H_k$) and there is no hyperplane of direction $\overline{H}_k$ and maximal weight which separates A_0 and $v_1 A_0$. Therefore

$$c_{y,v_1} \leq \sum_{i=m+1}^{i=n} c_{y,v_1}(\overline{H}_i) < \sum_{\substack{i=m+1 \\ i \neq k}}^{i=n} c_{y,v_1}(\overline{H}_i) + c_{\overline{H}_k} \leq \sum_{i=m+1}^{i=n} L(s_i) = L(w_{v_1}),$$

as required. $\square$

5.3. Proof of Theorem 4.3. In this section we fix $\lambda \in T$ and $z \in M_\lambda$. We set $v = w_\lambda z$. The following argument is inspired by a paper of Geck [3].

Lemma 5.4. *The submodule $\mathcal{M} := \langle T_x C_v \mid x \in X_\lambda \rangle_{\mathcal{A}} \subset \mathcal{H}$ is a left ideal.*

Proof. Since $\mathcal{H}$ is generated by T_s for $s \in S$, it is enough to check that $T_s T_x C_v \in \mathcal{M}$ for $x \in X_\lambda$. According to Deodhar's lemma (see [4, Lemma 2.1.2]), there are three cases to consider

- (1) $sx \in X_\lambda$ and $\ell(sx) > \ell(x)$. Then $T_s T_x C_v = T_{sx} C_v \in \mathcal{M}$ as required.
- (2) $sx \in X_\lambda$ and $\ell(sx) < \ell(x)$. Then $T_s T_x C_v = T_{sx} C_v + (v_s - v_s^{-1}) T_x C_v \in \mathcal{M}$ as required.

- (3) $t := x^{-1}sx \in S_\lambda$. Then $\ell(sx) = \ell(x) + 1 = \ell(xt)$. Now, since $tv < v$, we have (see in [7, §5.5, Theorem 6.6.b])

$$T_t C_v = v^{L(t)} C_v.$$

Thus, we see that

$$T_s T_x C_v = T_{sx} C_v = T_{xt} C_v = T_x T_t C_v = v^{L(t)} T_x C_v$$

which is in $\mathcal{M}$ as required. $\square$

Note that $\{T_y C_v \mid y \in X_\lambda\}$ is an $\mathcal{A}$ -linearly independent subset of $\mathcal{H}$. Indeed, for $y \in X_\lambda$ we have

$$\begin{aligned} T_y C_v &= T_y T_v + \sum_{u < v} P_{u,v} T_y T_u \\ &= T_{yv} + \text{an } \mathcal{A}\text{-linear combination of } T_w \text{ with } \ell(w) < \ell(yv). \end{aligned}$$

So, by an easy induction on the length and since the T_z form a basis of $\mathcal{H}$, we get the result. It follows that $\{T_y C_v \mid y \in X_\lambda\}$ is a basis of $\mathcal{M}$. Following [3], we shall now exhibit another basis of $\mathcal{M}$.

Lemma 5.5. *Let $y \in X_\lambda$, we can write uniquely*

$$T_{y^{-1}}^{-1} C_v = \sum_{x \in X_\lambda} \bar{r}_{x,y} T_x C_v, \quad \bar{r}_{x,y} \in \mathcal{A},$$

where $r_{y,y} = 1$ and $r_{x,y} = 0$ unless $x \leq y$.

Proof. We have

$$T_{y^{-1}}^{-1} = T_y + \sum_{z < y} \bar{R}_{z,y} T_z.$$

where $R_{y,v} \in \mathcal{A}$ (see [7, §4]). Now let $z \in W$ be such that T_z occurs in the above expression. We can write z uniquely under the form $x.w$ with $w \in W_\lambda$ and $x \in X_\lambda$ (see [4, Proposition 2.1.1]). Note that $x \leq z < y$. Since $xw = x.w$, we have $T_z = T_x T_w$ and

$$T_{y^{-1}}^{-1} C_v = T_y C_v + \text{an } \mathcal{A}\text{-linear combination of } T_x T_w C_v, \text{ where } x < y$$

Since $w \in W_\lambda$, we know that $\ell(w w_\lambda) = \ell(w_\lambda) - \ell(w)$ and $T_w C_v = v^{L(w)} C_v$. Therefore we see that $r_{y,y} = 1$ and $r_{x,y} = 0$ unless $x \leq y$. $\square$

Lemma 5.6. *Let $x, y \in X_\lambda$. We have*

$$\sum_{\substack{z \in X_\lambda \\ x \leq z \leq y}} \bar{r}_{x,z} r_{z,y} = \delta_{x,y}.$$

Proof. The proof is similar to the one in [3], once we know that $\{T_y C_v \mid y \in X_\lambda\}$ is an $\mathcal{A}$ -linearly independent subset of $\mathcal{H}$. $\square$

Proposition 5.7. *For any $y \in X_\lambda$, we have*

$$C_{yv} = T_y C_v + \sum_{x \in X_\lambda, x < y} p_{x,y}^* T_x C_v \quad \text{where } p_{x,y}^* \in \mathcal{A}_{<0}.$$

Proof. Fix $y \in X_\lambda$ and consider a linear combination

$$\tilde{C}_{yv} := \sum_{x \in X_\lambda, x \leq y} p_{x,y}^* T_x C_v$$

where $p_{y,y}^* = 1$ and $p_{x,y}^* \in \mathcal{A}_{<0}$ if $x < y$.

Our first task is to show that the p^* -polynomials can be chosen such that $\tilde{C}_{yv} = \overline{\tilde{C}_{yv}}$. In order to do so, we proceed as in the proof of the existence of the Kazhdan-Lusztig basis in [6]. We set up a system of equations with unknown the p^* -polynomials and then use an inductive argument to show that this system has a unique solution.

As in [3], one sees that the condition $\overline{\tilde{C}_{yv}} = \tilde{C}_{yv}$ is equivalent to

$$\sum_{x \in X_\lambda, z \leq x \leq y} \overline{p_{x,y}^*} \overline{r_{z,x}} = p_{z,y}^* \quad \text{for all } z \in X_\lambda \text{ such that } z \leq y.$$

In other words, the coefficient $p_{x,y}^*$ must satisfy

$$p_{y,y}^* = 1, \tag{1}$$

$$\overline{p_{x,y}^*} - p_{x,y}^* = \sum_{\substack{z \in X_\lambda \\ x < z \leq y}} r_{x,z} p_{z,y}^* \quad \text{for } x \in X_\lambda, x < y. \tag{2}$$

We now consider (1) and (2) as a system of equations with unknown $p_{x,y}^*$ ($x \in X_\lambda$). We solve it by induction. Let $x \in X_\lambda$ be such that $x \leq y$. If $x = y$, we set $p_{y,y}^* = 1$, so (1) holds.

Now assume that $x < y$. Then, by induction, $p_{z,y}^*$ ($z \in X_\lambda$) are known for all $x < z \leq y$ and they satisfy $p_{z,y}^* \in \mathcal{A}_{<0}$ if $z < y$. In other words, the right-hand side of (2) is known, we denote it by $a \in \mathcal{A}$.

Using Lemma 5.6 and the same argument as in [3] yields $\overline{a} = -a$. Thus the identity $\overline{p_{x,y}^*} - p_{x,y}^* = a$ together with the condition that $p_{x,y}^* \in \mathcal{A}_{<0}$ uniquely determine $p_{x,y}^*$.

We have proved that the coefficient $p_{x,y}^*$ can be chosen such that $\tilde{C}_{yv}$ is fixed by the involution $h \rightarrow \overline{h}$. Furthermore, we have

$$\begin{aligned} \tilde{C}_{yv} &= T_y C_v + \sum_{\substack{x < y \\ x \in X_\lambda}} p_{x,y}^* T_x C_v \\ &= T_y (T_v + \sum_{v_1 < v} P_{v_1,v} T_{v_1}) + \sum_{\substack{x < y \\ x \in X_\lambda}} p_{x,y}^* T_x (\sum_{v_1 \leq v} P_{v_1,v} T_{v_1}) \\ &= T_y T_v + \sum_{v_1 < v} P_{v_1,v} T_y T_{v_1} + \sum_{\substack{x < y \\ x \in X_\lambda}} \sum_{v_1 \leq v} p_{x,y}^* P_{v_1,v} T_x T_{v_1}. \end{aligned}$$

Now, in the above expression, the elements of the form $P_{v_1,v} T_x T_{v_1}$, with $x \in X_\lambda$ and $v_1 < v$, can give some coefficient in $\mathcal{A}_{\geq 0}$ (compare to the situation in [3]). However, by Lemma 5.2, we get

$$\tilde{C}_{yv} = T_{yv} + \text{an } \mathcal{A}_{<0}\text{-linear combination of } T_z \text{ with } \ell(z) < \ell(yv)$$

and then by definition of the Kazhdan-Lusztig basis, we conclude that $\tilde{C}_{yv} = C_{yv}$ as required. $\square$

Corollary 5.8. *We have*

$$\mathcal{M} = \langle C_{yv} \mid y \in X_\lambda \rangle_{\mathcal{A}}.$$

Proof. Let $y \in X_\lambda$. By Proposition 5.7 we have

$$C_{yv} = T_y C_v + \sum_{\substack{x \in X_\lambda \\ x < y}} p_{x,y}^* T_x C_v.$$

Thus $C_{yv} \in \mathcal{M}$. Now, by a straightforward induction, we see that

$$T_y C_v = C_{yv} + \text{a } \mathcal{A}\text{-linear combination of } C_{xv},$$

which yields the required assertion. $\square$

We can now prove Theorem 4.3.

Recall that

$$N_{\lambda,z} = \{w \in W \mid w = xv, x \in X_\lambda\}.$$

Let $y \in X_\lambda$. Let $z \in W$ such that $z \leq_L yv$. We want to show that $z \in N_{\lambda,z}$. To prove the theorem it is enough to consider the case where C_z appears with a non-zero coefficient in $T_s C_{yv}$ for some $s \in S$. By Corollary 5.8, $C_{yv} \in \mathcal{M}$. Since $\mathcal{M}$ is a left ideal, we know that $T_s C_{yv} \in \mathcal{M}$. Using Corollary 5.8 once more, we obtain

$$T_s C_{yv} = \sum_{y_1 \in X_\lambda} a_{y_1,yv} C_{y_1v}$$

and this expression is the expression of $T_s C_{yv}$ in the Kazhdan-Lusztig basis of $\mathcal{H}$. We assumed that C_z appears with non zero coefficient in that expression, therefore there exists $y_1 \in X_\lambda$ such that $z = y_1v$, and $z \in N_{\lambda,z}$ as required. The theorem follows.

REFERENCES

- [1] R. Bédard. The lowest two-sided cell for an affine Weyl group *Comm. Algebra* **16**, 1113-1132, 1988.
- [2] K. Bremke. On generalized cells in affine Weyl groups. *Journal of Algebra* **191**, 149-173, 1997.
- [3] M. Geck. On the induction of Kazhdan-Lusztig cells. *Bull. London Math. Soc.* **35**, 608-614, 2003.
- [4] M. Geck, G. Pfeiffer *Characters of finite Coxeter groups and Iwahori-Hecke algebras*, London Math. Soc. Monographs NS 21, Oxford University Press 2000.
- [5] G. Lusztig. Hecke algebras and Jantzen's generic decomposition patterns. *Advances in Mathematics* **37**, 121-164, 1980.
- [6] G. Lusztig. Left cells in Weyl groups. in *"Lie group representations", Lectures Notes in Math., Springer-Verlag, Vol. 1024, 99-111*, 1983.
- [7] G. Lusztig. Hecke algebras with unequal parameters. *CRM Monographs Ser.* **18**, Amer. Math. Soc., Providence, RI, 2003.
- [8] J.-Y. Shi. A two-sided cell in an affine Weyl group. *J. London Math. Soc. (2)* **36**, 407-420, 1987.
- [9] J.-Y. Shi. A two-sided cell in an affine Weyl group II. *J. London Math. Soc. (2)* **37**, 253-264, 1988.
- [10] N. Xi. Representations of affine Hecke algebras. *Lectures Notes in Math., Springer-Verlag, Vol. 1587*, 1994.

SCHOOL OF MATHEMATICS AND STATISTICS F07, UNIVERSITY OF SYDNEY NSW 2006, AUSTRALIA

E-mail address: guilhot@maths.usyd.edu.au